Agromet Advisory Bulletin for the District, Kannur





(Issued jointly by Kerala Agricultural University Regional Agricultural Research Station Pilicode& India Meteorological Department)



Bulletin Number:Pilicode/ Knr - 100/2025	Date: 16/12/2025

A. Weather Summary of preceding five days

Rainfall, mm	Max. temp., °C	Min. temp., °C	R. H., %	Wind speed, Km/h
0.0	34.6 – 35.4	20.6 – 222.0	52 – 65	06–12

B. Weather forecast for next five days

Parameters	17-12-2025	18-12-2025	19-12-2025	20-12-2025	21-12-2025
Average Rainfall, mm	0	0	0	0	0
Max. Temp, °C	34	34	34	34	34
Min. Temp,°C	21	21	21	21	21
Max. Relative Humidity, %	60	60	60	60	60
Min. Relative Humidity, %	55	55	55	55	55
Wind speed,km/h	3	4	3	3	3
Wind direction, degrees	270	270	270	270	270
Total cloud cover, octa	3	3	3	3	3

C. Agrometeorological Advisories

Cron	Stages	Problems	A cus matagual acidal advisauias	
Crop	Stages	riouleilis	Agro-meteorological advisories	
	No Rainfall** The sky will be partially cloudy. Atmospheric humidity will be normal. Low night ten and high day temperature will be experienced. Hence there will be distinct difference day time temperature and night temperature.			
General Condition				
Weather warning	No rainfall from December 17 to 21.			
Impacts	Damages and losses may occur for crops having slender stems like banana and vegetables.			
	High temperature during daytime may cause high rate of evaporation of water from the soil.			
	Direct exposure to sunlight may cause sunburn and injuries animals.			
General Recommendati	Give popping to all soft slender stemmed crops like banana, vegetables, climbers etc			
ons	free of any pest and di prevent the multiplicat soil using a secondary	seases. Avoid the trashes of the ion of crop specific pest and dis	the waterloss from soil. For mulching, the trashes used should be woid the trashes of the same species as mulch. This will help to specific pest and disease causing organisms. Powdering the top plement and spreading it uniformly over the field, will help to of time in the fields.	
	Adopt drip irrigation method for maximum water use efficiency.			
	Provide shade net for vegetable crops and ensure irrigation.			
	Donot keep animals under open conditions for a long time. Provide sufficient drinking water to them.			

Coconut	All stages	Basal rot	Drench the basin with Hexaconazole (25 ml in 25 L of water). After one week apply lime @ 2 kg per palm. Apply Potash @ 1 kg per palm three weeks after lime application to improve the health of palm, in addition to the regular dose. Avoid flood irrigation to prevent the spread of pathogen.	
Banana	Various stages of growth	Sigatoka leaf spot	Cut and burn all affected leaves. Spray Tilt or Contaf (@ 2ml per litre)	
Amaranthus	All stages	Leaf blight	Avoid overhead irrigation. Irrigate gently the basal regions of the plants. If disease is severe, apply cow dung slurry supernatant (2%) as foliar spray and Pseudomonas flourescens (2%) or Trichoderma viridae (2%) as soil application.	
Cucurbit Vegetables	Planting	Downy mildew	As a prophylactic measure apply 'Mancozeb'(@ 2g/l of water). If disease appeared, spray Akomin® (@3ml/L) or both surfaces of the leaves, thrice at 15 days interval. Remove and burn out the infested leaves.	
Cucurbitaceous Vegetables	Flowering and fruit setting	Melon fruit flies	Keep pheromone traps. Spray Malathion:- Mix 2 ml of Malathion in one litre of water. Dissolve 10g of jaggery in the solution to attract the flies. Spray this solution on lower sides of the leaves	
Chilli	Growth phase	Late spot	Spray Mancozeb or Saaf (2g/litre of water) or Potassium phosphonate (2ml/litre of water)	
Mango	Flowering stage	Mango hoppers	Spray Azadirachtin (@5ml/litre Azadirachtin 3000pmm)	

** Warning colour codes of rainfall (for disaster management)

Warning (Take actions)	Alert (Be prepared)	Watch (Be updated)	No warning (No actions)