



Agromet Advisory Bulletin for PALAKKAD District

(Valid from 22.03.2025 to 26.03.2025)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 12/2025, (23) Friday

Dated 21.03.2025

A. Medium range weather forecast of IMD for next five days

Weather parameters	22.03.2025	23.03.2025	24.03.2025	25.03.2025	26.03.2025
Rainfall (mm)	0.4	1	5	0.1	0.2
Maximum temperature (°C)	38	38	38	38	38
Minimum temperature (°C)	26	26	26	26	26
Maximum Relative Humidity (%)	90	90	90	90	90
Minimum Relative Humidity (%)	40	40	40	40	40
Wind speed (kmph)	4	3	4	3	2
Wind direction (deg)	160	70	200	20	110
Total cloud cover (octa)	8	8	8	7	6

Weather summary / Alert: -. As per the forecast of IMD, There is a chance of light rainfall in isolated places of the district. Above normal rainfall is expected in Kerala.

B. Agrometeorological Advisories for crop protection & management

General and Agrometeorological Advisories for Thrissur District

General Advisory: - As the climate is hot and dry, irrigate the crops as required to ensure adequate soil moisture. Avoid irrigation during strong sunlight. Drip irrigation can reduce water loss to some extent.

Crop	Pest/ Disease, Crop Management	Agromet Advisory
Puncha rice (Panicle Emergence stage)	Lakshmi disease	There is a chance of Lakshmi disease in rice. To control this, spray 2 gram Copper hydroxide in one liter of water.
	Brown Plant Hopper (BPH)	There is a chance of brown plant hopper attack in rice. Spray 2 gram Thiamethoxam per ten litre of water.
Cowpea	Thrips	There is a chance of thrips attack in cowpea. Apply 2% neem oil+ garlic emulsion under the leaf surface at 10 days intervals. If the attack is severe, apply spiromecifen 8 ml per ten litre of water.
Cucurbitaceous Vegetables	Mosaic disease	There is a chance of mosaic disease in cucurbitaceous vegetables. It can be controlled by rogue out the infected plants. Place yellow sticking traps in field to control the insect vector. If the attack is severe, spray Dimethoate 30 EC @ 1.5 ml per one litre of water.

Animal Husbandry: - Take precautions to avoid ecto-parasitic infestation. Fever, shivering, brown coloured urine, off feed are noticed, consult veterinary doctor immediately. (CAADECCS, KVASU, Mannuthy).