



Agromet Advisory Bulletin for PALAKKAD District

(Valid from 30.04.2025 to 04.05.2025)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara
based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 17/2025, (34) Tuesday

Dated 29.04.2025

A. Medium range weather forecast of IMD for next five days

Weather parameters	30.04.2025	01.05.2025	02.05.2025	03.05.2025	04.05.2025
Rainfall (mm)	5	5	3	2	2
Maximum temperature (°C)	38	38	38	38	38
Minimum temperature (°C)	25	25	25	25	25
Maximum Relative Humidity (%)	85	85	85	85	85
Minimum Relative Humidity (%)	65	65	65	65	65
Wind speed (kmph)	2	3	2	2	6
Wind direction (deg)	250	230	230	230	250
Total cloud cover (octa)	7	7	8	8	7

Weather summary / Alert: -. As per the forecast of IMD, there is a chance of light rainfall in the district. Thunderstorm accompanied by lightning and gusty winds is very likely to occur at one or two places over Kerala on 29th April to May 03rd 2025. Normal to above normal rainfall is expected in Kerala.

Maximum temperatures are very likely to be around 38 °C in 29th April to May 1st 2025. Hot & Humid weather is very likely to prevail over the district except in hilly areas due to high temperature and humidity.

B. Agrometeorological Advisories for crop protection & management

General and Agrometeorological Advisories for Palakkad District

General Advisory: - Proper mulching can be followed to conserve soil moisture. Since air temperature is increasing, it is a desirable practice to bury fresh or dried coconut husk in the coconut basin to conserve soil moisture.

Crop	Pest/ Disease, Crop Management	Agromet Advisory
Puncha rice (Maturity stage)	Brown Plant Hopper (BPH)	There is a chance of brown plant hopper attack in rice. Spray 2 gram Thiamethoxam per ten litre of water.
Rice (virippu)	Management	It is recommended to sow dhaincha seeds in the paddy field before virippu planting.
Banana	Calcium & Boron Deficiency	To control leaf curling in banana, due to calcium deficiency, apply 250g lime per banana plant. Boron deficiency is seen more during dry weather. Spray one gram borax per one litre of water.
Ginger, Turmeric	Planting time	While preparing land for ginger and turmeric cultivations, apply 2 to 3 kg lime per cent. As a prophylactic measure against rhizome rot disease, the ginger and turmeric rhizome should be soaked in 0.3% mancozeb or pseudomonas solution (3 gram per one litre of water) for 30 minutes.
Vegetables	Management	Addition of pseudomonas culture to the soil at the time of planting of vegetables will help to reduce fungal diseases

Animal Husbandry: - Before the rainy season begins, keep the cattle shed clean and tidy. (CAADECCS, KVASU, Mannuthy).