





Agromet Advisory Bulletin for Thrissur District

(Valid from 29.04.2025 to 04.05.2025)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 17/2025, (34) Tuesday

Dated 29.04.2025

			uring last to 29 th Ap)		Weather Parameters		Weather Forecast Valid for 30 th April to 04 th May 2025				
23/04	24/04	25/04	26/04	27/04	28/04	29/04	Date		30/04	01/05	02/05	03/05	04/05
12.7	0.0	0.0	0.0	0.0	0.0	18.3	Rainfall (mm)		4	4	2	1	0.5
34.9	33.8	34.4	34.5	35.1	36.3	35.6	$T_{max}(^{0}C)$		36	36	36	36	36
24.1	26.0	25.2	25.6	25.7	25.4	24.4	$T_{min}(^{0}C)$		24	24	24	24	24
Cloudy	Overcast	Cloudy	Cloudy	Cloudy	Cloudy		AM	Cloud	Cloudy	Cloudy	Cloudy	Cloudy	Cloudy
Cloudy	Partly cloudy	Cloudy	Cloudy	Cloudy	Cloudy		PM	cover					
98	95	87	93	92	94	95	RH-I (%)		85	85	85	85	85
73	67	59	71	66	70		RH-II (%)		75	75	75	75	75
2.5	2.5	2.0	2.2	2.3	2.8	3.2	Wind Spe	ed (km/hr.)	8	8	2	8	2
C	C	C	C	C	C	NE	AM	Wind	sw	wsw	sw	sw	wsw
SSW	SSW	NW	SSW	SSW	SW		PM	direction					
Rainfall (mm) in last week R				Rainf	Rainfall (mm) from 01/01/2025 to till dated				Rainfall (mm) from 01/03/2025 to today				
31.0				124.9					124.9				

Weather summary / Alert: -. As per the forecast of IMD, there is a chance of light rainfall in the district. Thunderstorm accompanied by lightning and gusty winds is very likely to occur at one or two places over Kerala on 29th April to May 03rd 2025. Normal to above normal rainfall is expected in Kerala.

Maximum temperatures are very likely to be around 36°C in 29th April to May 1st 2025. Hot & Humid weather is very likely to prevail over the district except in hilly areas due to high temperature and humidity.

General and Agrometeorological Advisories for Thrissur District

General Advisory: - Proper mulching can be followed to conserve soil moisture. Since air temperature is increasing, it is a desirable practice to bury fresh or dried coconut husk in the coconut basin to conserve soil moisture

Crop	Pest/ Disease, Crop Management	Agromet Advisory				
Puncha rice	BrownPlant	There is a chance of brown plant hopper attack in rice. Spray 2 gram				
(Maturity stage)	Hopper(BPH)	Thiamethoxam per ten litre of water.				
Rice (virippu)	Management	It is recommended to sow dhaincha seeds in the paddy field before virippu planting.				
	Calcium & Boron	To control leaf curling in banana, due to calcium deficiency, apply				
Banana	Deficiency	250g lime per banana plant. Boron deficiency is seen more during				
	Betterency	dry weather. Spray one gram borax per one litre of water.				
		While preparing land for ginger and turmeric cultivations, apply 2 to				
		3 kg lime per cent. As a prophylactic measure against rhizome rot				
Ginger, Turmeric	Planting time	disease, the ginger and turmeric rhizome should be socked in 0.3%				
		mancozeb or pseudomonas solution (3 gram per one litter of water)				
		for 30 minutes.				
Vegetables	Management	Addition of pseudomonas culture to the soil at the time of planting of				
vegetables	wianagement	vegetables will help to reduce fungal diseases				
Animal Husbandry: - Before the rainy season begins, keep the cattle shed clean and tidy. (CAADECCS, KVASU, Mannuthy).						