



Agromet Advisory Bulletin for Thrissur District

(Valid from 18.05.2024 to 22.05.2024)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara
based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 20/2024, (40) Friday

Dated 17.05.2024

Observed weather during last week (Dated 11 th May to 17 th May 2024)							Weather Parameters		Weather Forecast Valid for 18 th May to 22 nd May 2024				
11/05	12/05	13/05	14/05	15/05	16/05	17/05	Date		18/05	19/05	20/05	21/05	22/05
0.6	25.1	0.0	3.3	6.6	0.0	1.9	Rainfall (mm)		5	4	12	23	23
34.2	35.9	35.4	32.2	32.3	36.2	36.3	T _{max} (⁰ C)		35	34	34	34	34
25.9	24.8	26.4	25.8	23.3	25.3	26.0	T _{min} (⁰ C)		26	25	25	25	25
Cloudy	Cloudy	Cloudy	Overcast	Cloudy	Cloudy	Cloudy	AM	Cloud cover	Cloudy	Cloudy	Cloudy	Cloudy	Cloudy
Overcast	Overcast	Cloudy	Cloudy	Overcast	Overcast		PM						
95	92	95	92	93	84	89	RH-I (%)		87	87	87	87	87
72	69	83	78	64	62		RH-II (%)		71	71	71	71	71
1.1	1.5	1.5	1.2	1.7	2.8	1.7	Wind Speed (km/hr.)		2	2	2	2	14
C	C	C	C	NNE	C	C	AM	Wind direction	SW	SW	SW	SW	SW
C	SW	ESE	C	SSW	WSW		PM						
Rainfall (mm) in last week				Rainfall (mm) from 01/01/2024 to till dated				Rainfall (mm) from 01/03/2024 to today					
37.5				191.9				59.7					

Weather summary / Alert: Thunderstorm with lightning accompanied by gusty wind speed reaching 30 to 50 Kmph is very likely to occur at one or two places in Kerala from 17th to 21st May 2024.

SMS: - As per the forecast of IMD, there is a chance of Moderate rainfall (15.6 to 64.4mm) is expected in isolated places of the district. Above normal rainfall is expected in Kerala.

General and Agrometeorological Advisories for Thrissur District

Special crop management is required for agricultural crops before monsoon season. Proper drainage facilities have to be ensured. Provide supports for the bananas. Make the vegetable pandals stronger. Avoid standing in open areas to protect farmers and livestock from lightning and thunder. Proper mulching can be followed to conserve soil moisture.

Crop	Pest/ Disease	Crop/ Animal Advisory
Rice (virippu)	-	It is recommended to sow dhaincha seeds in the paddy field before virippu planting. Land preparation and nursery preparation for virippu paddy (Rainfed) can be started after receiving summer rains.
Coconut & Arecanut	-	Pits can be taken for planting seedlings of coconut and arecanut.
Ginger, Turmeric	Planting time	While preparing land for ginger and turmeric cultivations, apply 2 to 3 kg lime per cent. As a prophylactic measure against rhizome rot disease, the ginger and turmeric rhizome should be soaked in the 0.3 percent mancozeb or pseudomonas solution (3 gram per one litter of water) for 30 minutes. Moreover 90 kg FYM mixed with one kg Trichoderma and kept under shade for two weeks with sufficient moisture and can be applied.

Animal Husbandry: - Deworm the animals after dung examination. Care should be taken to prevent rat breeding in the premises of cattle shed to prevent leptospirosis. Keep away livestock from open fields to avoid lightning and thunder.